ENVS 491
Senior Seminar
Fall 2003

Dena Bober
Brandi Hale
Michael Hansen
Jennelle Kershaw
Timothy Marshall
Julie Okronley
Matt Rescott

Table of Contents

Acknowledgements iii

Preface iv

I. Introduction 1
Ephraim Maduabuchi
Jenelle Kershaw

II. History 4
Erin Shutt

III. Current Practices 7
Mike Malone

IV. Impacts of Landscape 9
Human Aspects 9
Tim Marshall
Natural Aspects 11
Mike Hansen
Elizabeth Yarbrough
Brandi Hale
Jennelle Kershaw

V. Landscape Stewardship 16
Rain Gardens 16
Matt Rescott
Riparian Buffer Zones 17
Jennelle Kershaw
Impervious vs. Pervious Surfaces 19
Julie Okronley
BayScaping 19
Elizabeth Yarbrough
Cow Paths 20
Mike Malone

VI. Towson As a Model 22
Mike Hansen
Elizabeth Yarbrough
Curt Kleinsorg
Tim Marshall

VII. New Initiatives 25
Erin Shutt
Matt Rescott

VIII. Conclusion 28
Dena Bober

IX. References Cited 29

X. Appendices 32
Brandi Hale
Jennelle Kershaw

Acknowledgements

We would like to acknowledge the following people for their kind assistance during this semester. We hope that we properly used the information that you provided us.

Mr. Paul Thomas -- Landscape Services Manager at Towson University

Mr. Steve Saari -- Watershed Coordinator, Interstate Commission on the Potomac Basin

Ms. Jeanne Armacost -- DEPRM, Natural Resource Specialist

Ms. Nancy Gonce -- Library archivist at Towson University

Ms. Ada Woods -- Reference Librarian at Towson University

Dr. Martin Roberge -- Geography Professor at Towson University

Dr. Jon Filer -- Geology Professor at Towson University

Ms. Valerie Gaine -- United States Geological Survey

Dr. Jane Wolfson -- Director, Environmental Science and Studies

Preface

All students completing the Environmental Science and Studies major enroll in ENVS 491, Senior Seminar, during their senior year. In this course, students are presented with

developing solutions/suggestions that are economically sound, logistically feasible and that incorporate stakeholder needs and constraints.

I. Introduction

The landscape of Towson University is the front door of a complex academic institution. It plays important roles in education, recruitment, recreation, and university

are presented. In addition, ideas to reduce the impacts of the student population on the landscape will be presented.

As an integral part of the Towson community, Towson University can become a model for other institutions and communities interested in a more ecological approach to landscaping. Several suggestions of how to enhance current practices on campus creating a more ecosystem-based approach will be a focus.

II. History

recognition of University history is essential. Towson University, known previously by several different names, did not originate in the town of Towson, Maryland.

Over the past 135 years, Towson University has undergone some major changes, not only educationally, but aesthetically as well. In 1865, the State legislature allocated -training school. In 1866, The State Normal School opened on rented land in Baltimore City. M.A. Newell, the first principal, oversaw 11 students and 3 faculty members (N. Gonce, personal communication, 2003). After occupying two rented sites in Baltimore City between 1866 and 1876, the 10 faculty members and 206 students of the State Normal School moved. The first building constructed specifically for the school was located at the corner of Carrollton and Lafayette Avenues in Baltimore City (N. Gonce, personal communication, 2003).

On April 24, 1904, John and Mary Nelligan purchased 5 plots (72 acres) of land that would eventually be known as Towson University. The 5 plots were bought from Richard S. Albert, Abraham P. Nonemaker, Ray P. Allen, Arthur L. Bosley, and Richard Pleasant, who were trustees of the land at the time of purchase. The true landowner is not

personal communication, 2003). The new State Normal School campus, comprising the Administration Building (now Stephens Hall), Newell Hall, and the Power Plant, was dedicated

new demands, and by 1976, the college was renamed Towson State University (N. Gonce, personal communication, 2003).

In 1988, the university became a part of the University System of Maryland. Yet again in 1997, the college was renamed Towson University to reflect the evolution from a State-supported to a State-assisted institution. Towson University has evolved from a small school in downtown

higher education. We now have 17,000 undergraduate and graduate students enrolled in 90 educational programs encompassing liberal arts and sciences, as well as applied professional fields (N. Gonce, personal communication, 2003).

Paul Thomas is responsible for both the maintenance and aesthetics of the campus landscape. His crew is an integral part of connecting the institution and its campus to the surrounding community. An improvement of the current practices can only further

IV. Impacts of Landscape

Human Aspects

students, and surrounding communities. Knowing how different landscapes affects the mental behavior of the inhabitants is important and can help us develop an effective landscape design.

The appearance of trash on the landscape may have negative affects upon human health. The visual distraction created by trash is found throughout campus from the woods to the classroom. Such distraction causes people to be indirectly affected by their physical environment. A well-maintained landscape with features that increase the aesthetics of the area also influences the way people are likely to treat it. When students begin to valu

prevalent. Random littering by the student body may decrease and students throughout the institution may make the effort to help keep campus clean. Such actions will decrease the costs incurred by maintenance and will also provide a more visually appealing campus.

mental approach. The stresses associated with urbanized areas, such as noise, crowding, pollution, and traffic, are domineering factors in the mental approach to everyday living (Kuo et al., 1998). While some people may feel comfortable and work well in such environments, others may not. Depending on one's point of view, a rural area or natural landscape may be preferred for its tranquility, beauty, and natural features (Hall IV et al.,

1989). It has been shown in qualitative studies that a pleasant landscape can change or equalize the mood of inhabitants (Shang et al., 2000). While an urban setting offers little

promote a sense of tranquility and offer some natural healing properties (Staats et al., 2003).

Although most may not know what kind of tree or grass they are looking at, people make instant associations with what they see (Purcell et al., 2001). Some people also feel safer in an urban environment that includes well-placed vegetation. Vegetation cannot be so dense as to limit the ability that one has to observe their surroundings (Herzog et al., 2000). By providing an adequate amount of properly placed vegetation, feelings of safety may be improved. Easing the minds of students and faculty through proper landscape design may be a good way to decrease a fear that some may feel.

Moreover, landscaped areas allow for better recovery from fatigue (Staats et al.

Path choice is also influenced by the present landscape. Pedestrians often choose the shortest and easiest path. Paths that are littered with signs, restrictions, and have a bland appearance are generally avoided (Zacharias, 2001). Path dynamics on Towson Univers

Providing interesting path options in appropriate directions could eliminate undesirable

of its infrastructure. It is essential that Towson University consciously consider hydrology since actions taken on campus directly affect surrounding communities through its flow off campus.

There are two main streams on campus, South Fork Towson Run and main branch Towson Run. The two streams eventually meet and drain into Lake Roland. The origin of these streams is off campus east of York road. Due to the substantial development of the area, some streams have been completely translated into pipe and put underground. Such is the case with much of the Glen Tributary, a small stream on campus in the Glen, which drains into Towson Run. The first time the water in the Glen encounters daylight is when it enters onto campus in its newly restored channel.

The geology department, under the initiative of Dr. Filer, has begun a small monitoring network of campus hydrology. This network includes 2 stream gauges, a groundwater well, a rain gauge, and 2 sites to sample water quality. It is important to monitor hydrology, as any landscape management plan will have an impact. This network begins to describe the hydrology of campus. An expansion of this network will lead to a better understanding of campus hydrology.

The campus landscape provides unique educational opportunities as the

ting creates unique effects on streams. Urban settings induce unnatural flow characteristics in terms of supply to stream channels. Unnatural flow characteristics may range from frequency of flood events to simply an increase in suspended sediment.

An understanding of the current properties of the water in streams must be applied

Faculty and students could implement and maintain an impressive hydrologic monitoring network. Stations for stage and discharge of streams could be expanded to include other parts of campus and further understand the journey of water. Groundwater wells could be used to monitor changes in water quality and illustrate effects of

Control of invasive species, which occupy areas between different land uses, should occur. Filling these areas with native natural landscape elements could do this. Wildflower beds and buffer plants are excellent in this role. By using such natural methods, stewardship for the landscape by its users will increase.

V. Landscape Stewardship

Several methods can be used to facilitate environmental stewardship, which are well-developed innovative solutions. Modification of our landscape to include such

environment.

Rain Gardens

A need exists for innovative methods to control storm water runoff during and

effective at reducing the energy associated with runoff. Low areas that allow water to pool can be lined with grass, mulch or gravel.

Rain gardens require little maintenance upon completion especially when compared to lawns. Like all gardens, rain gardens require routine upkeep to continue maximum benefit. Their bioretention properties, and simple maintenance needs ensure an ecological storm water control tool.

Riparian Buffer Zones

Riparian zones are commonly defined as vegetative corridors adjacent to stream channels. Flooding, erosion, pollution and habitat degradation are all problems associated with the waterways on campus. Increasing the width of the vegetated riparian zones can significantly affect such occurrences by providing a barrier, slowing runoff and filtering water prior to discharge into the stream.

The size of the buffer needed depends on the size of the stream or frequency of flooding. Essentially the larger the width of the buffer the greater the volume it can filter runoff. Well-maintained riparian zones are at least 100 feet on each side of the stream

with an environmentally safe solution that hardens the soil. Though the soil becomes a hardened surface, pores still remain allowing for infiltration. As these innovative surfaces increase across campus, surface temperatures during the day will show a slight decrease. That decrease is the direct result of a change in surface color. Lighter surface colors reflect solar radiation and therefore discourage heat absorption. Shading from trees will also contribute to lower surface temperatures (NEMO, 2003).

While the absorption of heat is of concern to the campus community, so is the performance of a surface in colder conditions. Snow is particularly difficult to handle when working with alternative surfaces. Research suggests that more porous surfaces are able to thaw at a faster rate. With a rubber blade placed on the plow, snow removal is no longer a problem (NEMO, 2003).

As Towson's campus continues to grow, it is important to approach developmental issues with landscape stewardship in mind. By increasing the amount of permeable surfaces, the quantity and strength of storm water runoff produced on campus will decrease. Taking new initiatives to improve infiltration will increase the water quality of local streams and provide greater protection against soil erosion. Undisturbed soil is needed to store moisture and nutrients in order to sustain healthy vegetation.

BayScapes

BayScaping is a new and innovative way to landscape that was developed to prevent further harm to the Chesapeake Bay. BayScapes are environmentally sound and have many benefits for wildlife and people. Less mowing and fertilizer applications are required in areas in which they have been installed. These landscapes are colorful and help protect water quality. BayScapes also provide habitats for songbirds, small

mammals, butterflies, and a plethora of other organisms. BayScaping is being promoted throughout the Chesapeake Bay watershed and people are encouraged to visit representative site (ACB 2003).

BayScaping on Towson University's Campus could greatly improve Towson

storm events expose underlying rock. Conditions on the path worsen and cause it to become an undesirable place to walk. Pedestrians will now choose to walk along the sides of the path because it is flatter and easier to navigate. This, in turn, kills more grass and the path is continually widened (DNP, 2003).

As noted earlier, creating attractive features that draw pedestrian traffic can be a good way to influence path choice. Additionally, roping off specific areas or planting obstacles, such as trees and shrubs, can decrease

VI. Towson University as a Model

By adopting environmental stewardship as a long-term goal, and implementing programs to attain this goal, Towson has the potential to expand its role as an institution of learning. As environmental awareness continues to build within our culture, it is important that institutions of all kinds step forward to set an example for others to follow. A model is an example or a demonstration of a concept or idea. As a model we could serve to illustrate the ideas of environmental stewardship to those who want to see what can be done. A model can be applied to multiple cases and specialized to fit certain situations. By initiating environmental stewardship in its landscape and education program, Towson University becomes a mechanism by which other people and organizations could increase their environmental responsibility. If Towson were to become a model institution, the university stands to benefit in several ways.

The current campus landscape, potential initiatives that could be implemented, and some technical information have all been addressed thus far. While this is important, it is equally important to address the reasons why such actions should be taken. As we implement programs to make our landscape more environmentally responsible and friendly, we assume the role of a model. Since the level of ecological awareness we would be demonstrating is not prevalent, the university becomes the exception rather than the norm. There are two important groups of people for whom the university will serve as a model, residents of the surrounding community, and other institutions. Towson is situated in a unique niche where it can serve as a model for both.

The University has indicated it plans to become more involved in the surrounding community in the next few years and the campus landscape is an ideal place to start.

Primarily because of its visibility, the campus landscape provides an arena in which the university can show community constituents the innovative changes as they are implemented on campus. Residents who see these changes might want to adapt those ideas to the landscapes in which they reside or at the institutions at which they work. Our

go out into the community to share what is has been doing with those who may not be familiar with the campus. Towson University can take a proactive approach to its involvement in the community. One means might be to become further involved in the Jones Fall Watershed association which is our local watershed (Towson Run flows into Jones Falls). Another option might be to start a watershed association specifically for Towson Run. In this venue, Towson University can demonstrate how the parts of its landscape management program benefits the stream. Watershed associations are popular in the Baltimore region with the development of three major organizations and smaller spin offs in local areas. For example the Bear Creek Watershed Association in Dundalk is a spin off of the Herring Run Watershed Association.

Realizing the value of becoming a model campus, several universities already have management plans that will make them environmentally sustainable. Many Ivy League schools make a reference to environmental stewardship in their Master Plan and statement of university mission. For example, Brown University's "Brown is Green"

Is Green initiative in 1990 to facilitate the conservation of resources, waste reduction

University, 2003). Cornell University has also made its environmental mission well

known. The head of the Center for the Environment represents a group of alumni that sponsors programs for discussion and networking between students, faculty and Council members actively addressing environmental issues (Cornell University, 2003). Not only Ivy League schools are declaring their intentions regarding environmental responsibility.

conscious students to their institutions. Implementation of our new master plan provides a golden opportunity to actively deploy and advertise our environmental stewardship. If Towson University were to join the growing list of colleges that are known for taking care of the environment, we will improve our reputation at both the national and local levels.

The area west of Osler Drive could be developed as a recreation and relaxation area. A restoration proposal developed by Dr. Hull and colleagues in the Biology Department presents a plan for the use of this area (Wolfson, personal communication, 2003). This area can be incorporated into several classes. Ecology, limnology, and herpetology classes could benefit from the use of the proposed ponds and wetland. Restoration of this area could result in an extraordinary facility on campus.

VIII. Conclusion

Our project required that we, the students, possessed a clear understanding of several concepts and ideas. These concepts and ideas are what form the basis for our suggestions to campus planners. The following is our sense of what some terms mean. Landscape elements combine to form landscapes. A landscape is a mosaic, or a cluster of interacting ecosystems that is repeated throughout a specific location, no bigger than about 10,000km^2 (Huggett, 2000). Ecosystems are a web of intricately complicated life forms each relying on the other for survival. The purpose of this entire project has been to inform and educate those who read it. It has been our intent to convey our sense of urgency to other universities, communities, organizations, etc. We hope to initiate a chain reaction started from our example. We hope that information we provide about the potential for the future of campus results in implementation of some of our suggestions.

If we consider the Towson University campus landscape as just one puzzle piece in the big picture, then by enriching our environmental stewardship, we complete our portion of the puzzle and contribute to regional environmental improvement.

Our intentions have been to: (1) address the specific environmental challenges

Kuo F.E, Magdelena B. 1998. Transforming inner-city landscapes: Trees, sense of safety, and preference. Environment and Behavior 30: 28-59.

(LDNPA) Lake District National Park Authority. 2003. Educational Service: Footpath Erosion Factsheet. http://www.glowingcoast.co.uk/lakedist/access/info/erosion.pdf. (25 Oct 2003).

(NEMO) Nonpoint Education for Municipal Officials. 2003. Reducing Runoff. University of Connecticut. http://nemo.uconn.edu/reducing_runoff.html. (9 Oct 2003).

(NPS) National Park Service. 2001. Native Plants for Wildlife Habitat and Conservation Landscaping in Maryland. http://www.nps.gov/plant/pubs/nativesMD/info.htm. (22 Sept 2003).

(NPS) National Park Service. 2002. Plant Invaders of Mid-Atlantic Natural Areas.

Nye, J. 2003. Personal Communication. Director of Planning, Facilities Management, Towson University.

Petrides G, Wher J. 1998. Eastern Trees. New York: Houghton Mifflin Company.

(PGDER) Prince Georges County Department of Environmental Resources. 2003. Rain Gardens: The natural solution.

Purcell T, Peron E. 2001. Why do preferences differ between scene types. Environment and Behavior 33: 93-106.

Seedland. 2003. Growing a Beautiful Tomorrow. http://www.seedland.com (17 Oct 2003).

Shang H, Bishop I.D. 2000. Visual thresholds for detection, recognition and visual impact in landscape settings. Journal of Environmental Psychology 20: 125-140.

Smittle, D. 2002. Care-free plants. Pleasantville, N.Y. :Readers Digest. Association Inc.

Staats H, Kieviet A. 2003. Where to recover from attention fatigue: An expectancy- value analysis of environmental preference. Journal of Environmental Psychology 23: 147-157.

Thomas, P. 2003. Personal communication. Manager Landscape Services, Towson University.

Thompson, L. 1999. Native Plants for Central Maryland Landscape: Maryland Flora. http://www.mdflora.org/publications/natplants.html

Appendices

Appendix A
Recommendations for Control and Replacement.

Brandi Hale and Jennelle Kershaw
(Information presented below obtained from Westbrooks, 1998)

Norway Maple *Acer platanoides*

This tree has large, five-point leaves similar to sugar maple. When the leaf or stalk is broken a drop of white sap will show. The fall foliage is yellow. This tree has allelopathic properties preventing the growth of grass, garden plants, and forest understory beneath it. The seeds travel in the wind and can grow in deep shade. Without control the Norway maple could take over an entire forest. To control this tree, seedlings can be pulled by hand and large trees can be cut down. Herbicides can also be used to control the Norway maple but is not highly recommended. Some native alternatives include: American beech, Red maple, and Willow Oak.

Tree of Heaven *Ailanthus altissma*

This tree has long, compound leaves, with 11-25 lance-shaped leaflets that smell like peanut butter or burnt coffee when crushed. It is incredibly virulent and can grow under very poor conditions. It is a major threat because it produces a huge number of wind-borne seeds. It also grows very quickly and has allelopathic properties that kill surrounding plants. To control this invasive tree, seedlings can be pulled up but one should make certain to remove the entire plant and all of its root structure, if not the tree will grow back.

production. Herbicides is also an effective control but not highly recommended. Some native alternatives include: Wild ginger, Lady fern, and Creeping Phlox.

Mile-A-thumb *Polygonum perfoliatum*

This annual non-native weed has triangular leaves, barbed stems, and turquoise berries. It rapidly grows on slopes covering and shading out surrounding plants. Controlling this vine involves pulling seedlings and vines out by hands before seed set. Herbicides are also an effective means to controlling Mile-A-Minute.

Appendix B -

Brandi Hale and Jennelle Kershaw
(Information obtained from Hawthorne, 1999 and Smittle, 2002)

Shade Tolerant Plants:

Bleeding Heart

Appendix D Trees Suitable for the Towson University Campus

Joe Vogelpohl
(Information below obtained from Dirr, 1983; DNR, 2003; NPS, 2001, 2002; Petrides & Wher, 1998; and Thompson, 1999)

Trees are listed according to their preference of soil moisture and height.

Red Maple (*Acer rubrum*)

The Red Maple is probably the most adaptable tree you can plant. It grows well in wetland areas including streams but also can be planted in the upland areas. Do not plant it near any sidewalks or anywhere its roots can push up causing problems. The tree has very strong wood and nice fall colors. Typically the tree

Green Ash

Fringetree (*Chionanthus virginicus*)

The Fringetree likes deep moist fertile soil but can grow on dryer upland sites. This tree grows normally between 25-
rate is slow just about under a foot a year. The tree prefers an acidic soil of 6.0 to 6.5. It likes full sun on stream banks and grows well in clusters. Also it does very well in cities and has a high tolerance to air pollution. The tree has a half to two thirds an inch dark blue drupes (fruit), which birds love. This tree also produces large clusters of aromatic white flowers in late spring.

White Oak (*Quercus alba*)

The White Oak grows best in moist woodland sites that are well drained. This tree is typically between 60-

cannot tolerate poorly drained soil, or particularly dry soil, and the tree can get uprooted if winds are severe. The American Holly can tolerate higher air pollution and high soil pH. This tree is an evergreen

Chestnut Oak (*Quercus prinus*)

This tree does best in well-drained soils and does well in rocky dry and poor upland sites. This tree is a dominant tree around here. The normal size is from 60

Pignut Hickory (*Carya glabra*)

The Pignut Hickory will grow on very rocky dryer soil. It grows between 80-

cause problems in clean up, damage to cars, and could cause a student quite a lot of pain. Do not suggest planning this tree in areas where it could cause these problems.

Scarlet Oak (*Quercus coccinea*)

The Scarlet Oak is similar to the other Oaks but it likes dry sandy to rocky soil. The tree grows between 70-7
to two feet a year.

Post Oak (*Quercus stellata*)

The Post Oak is similar to other Oaks; it likes rocky ridges and silty loams but can grow in dry gravelly or sandy soils. This is a smaller Oak which grows typically between 40-